



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the posts, etc., outside the car seem passing by on account that we judge mostly by what is near our eyes. It is this that being stationary with respect to the earth, the ancients imagined the sun and stars to move around the earth instead of the opposite.

GEOMETRY.

110. Proposed by P. S. BERG, A. M., Principal of Schools, Larimore, N. D.

If the three face angles of the vertical triedral angle of a tetraedron are right angles, and the lengths of the lateral edges are represented by a , b , and c , and of the altitude by p , then $1/p^2 = 1/a^2 + 1/b^2 + 1/c^2$. [Chauvenet's *Geometry*.]

I. Solution by GEORGE E. DEAN, Professor of Mathematics, School of Mines and Metallurgy, University of Missouri, Rolla, Mo.; P. H. PHILBRICK, C. E., Lake Charles, La.; and CHAS. C. CROSS, Libertytown, Md.

Let OC , OB , OA be the edges mutually at right angles; OP the perpendicular. Join AP and produce to meet CB at D ; join BP and produce to meet AC at E ; join CP and produce to meet AB at F .

$$\text{Then } \frac{\triangle APC}{\triangle ABC} + \frac{\triangle BPC}{\triangle ABC} + \frac{\triangle APB}{\triangle ABC} = 1.$$

$$\text{But } \frac{\triangle PAC}{\triangle ABC} = \frac{PE}{BE}, \quad \frac{\triangle BPC}{\triangle ABC} = \frac{DP}{AD},$$

$$\frac{\triangle BPA}{\triangle ABC} = \frac{PF}{CF}.$$


$$\text{Therefore } \frac{PE}{BE} + \frac{DP}{AD} + \frac{PF}{CF} = 1.$$

$$BE \times BP = OB^2 = b^2; \quad BE = \frac{b^2}{BP}; \quad \frac{PE}{BE} = \frac{PE \times BP}{b^2} = \frac{OP^2}{b^2}.$$

$$AD \times AP = OA^2 = a^2; \quad AD = \frac{a^2}{AP}; \quad \frac{DP}{AD} = \frac{AP \times DP}{a^2} = \frac{OP^2}{a^2}.$$

$$CF \times CP = OC^2 = c^2; \quad CF = \frac{c^2}{CP}; \quad \frac{PF}{CF} = \frac{PF \times CP}{c^2} = \frac{OP^2}{c^2}.$$

$$\text{Hence } \frac{OP^2}{a^2} + \frac{OP^2}{b^2} + \frac{OP^2}{c^2} = 1, \text{ or } \frac{1}{p^2} = \frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2}. \quad \text{Q. E. D.}$$



II. Solution by P. S. BERG, B.Sc., Superintendent of Schools, Larimore, N. D.; WALTER H. DRANE, Graduate Student, Harvard University, Cambridge, Mass.; ELMER SCHUYLER, High Bridge, N. J.; W. H. WILSON, Professor of Mathematics, Geneva College, Beaver Falls, Pa.; and G. B. M. ZERR, A. M., Ph.D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

Let $OP=p$, $OC=a$, $OA=b$, $OR=c$, $OI=l$, $IH=m$, $PH=n$.

The triangles (right angled) OPA and OPI are similar, having $\angle POA$ in common.

$$\therefore p/b=l/p, \text{ or } p^2/b^2=l^2/p^2, \text{ or } p^4/b^2=l^2.$$

$$\text{Similarly, } p^4/a^2=n^2, p^4/c^2=m^2.$$

$$\therefore p^4/a^2 + p^4/b^2 + p^4/c^2 = n^2 + l^2 + m^2 = p^2.$$

$$\therefore 1/a^2 + 1/b^2 + 1/c^2 = 1/p^2.$$

Also solved by J. SCHEFFER, and E. D. SCALES.

111. Proposed by GEORGE R. DEAN, Professor of Mathematics, University of Missouri School of Mines and Metallurgy, Rolla, Mo.

Given that the area of a triangle is equal to half the product of two sides and the sine of the included angle, prove that $\sin(x+y) = \sin x \cos y + \cos x \sin y$.

I. Solution by W. F. BRADBURY, A. M., Head Master, Cambridge Latin School, Cambridge, Mass., and B. F. YANNEY, A. M., Professor of Mathematics, Mount Union College, Alliance, O.

Given area $\triangle = \frac{1}{2}ac \sin B \dots (1)$.

$$\sin B = \sin[180^\circ - (A + C)] = \sin(A + C).$$

$$\therefore \text{Area } \triangle = \frac{1}{2}ac[\sin(A + C)] \dots (2).$$

Draw BD perpendicular to AC .

$$\text{Area } \triangle = \frac{1}{2}BD(AD + DC).$$

But $BD = c \sin A = a \sin C$, and $AD = c \cos A$, and $DC = a \cos C$.

$$\therefore \text{Area } \triangle = \frac{1}{2} \frac{c \sin A}{a \sin C} \text{ or } (c \cos A + a \cos C) = \frac{1}{2}(ac \sin C \cos A + ac \sin A \cos C) \dots (3).$$

Putting (2) = (3), $\frac{1}{2}ac[\sin(A + C)] = \frac{1}{2}ac(\sin A \cos C + \cos A \sin C)$ or $\sin(A + C) = \sin A \cos C + \cos A \sin C$.

II. Solution by J. OWEN MAHONEY, B. E., Professor of Mathematics and Science, Carthage High School, Carthage, Tex.; J. W. YOUNG, Columbus, O.; J. SCHEFFER, A. M., Hagerstown, Md.; E. L. SHERWOOD, A. M., Professor of Mathematics, Whitworth College, Miss.; WALTER H. DRANE, Graduate Student, Harvard University, Cambridge, Mass.; B. F. SINE, Principal of Normal School, Capon Bridge, W. Va.; G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.; JOHN MACHNIE, A. M., Professor of Latin, University of North Dakota; H. F. STRATTON, Student in Heidelberg University, Tiffin, O.; J. C. NAGLE, C. E., Professor of Civil Engineering and Physics in the Agricultural and Mechanical College of Texas, College Station, Tex.; CHARLES C. CROSS, Libertytown, Md.; and ELMER SCHUYLER, High Bridge, N. J.

PROOF. Consider the triangle ACB .

The area of $ACB = \frac{1}{2}ac \sin B = \frac{1}{2}ac \sin(A + C) = \frac{1}{2}hb = \frac{1}{2}h(AD + DC)$, or

$$\begin{aligned} \sin(A + C) &= \frac{h}{a} \cdot \frac{AD}{c} + \frac{h}{c} \cdot \frac{DC}{a} \\ &= \sin C \cos A + \sin A \cos C. \end{aligned}$$

$$\therefore \sin(x + y) = \sin x \cos y + \cos x \sin y.$$